arXiv:1401.1589v2 [math.AP] 9 Jan 2014

A SINGULAR INTEGRAL APPROACH TO THE MAXIMAL L^p REGULARITY OF PARABOLIC EQUATIONS

BUYANG LI

1 Introduction

Consider a parabolic problem

$$\partial_t u + Au = f \tag{1.1}$$

on a Banach space X. The maximal L^p regularity of the parabolic problem (1.1) refers to such estimates as

$$\|\partial_t u\|_{L^p(\mathbb{R}_+;X)} + \|Au\|_{L^p(\mathbb{R}_+;X)} \le C\|f\|_{L^p(\mathbb{R}_+;X)}, \quad \text{for } 1 < p < \infty. \tag{1.2}$$

Traditionally, this estimate was derived for parabolic equations by using parabolic singular integrals [1, 6] for $X = L^p$. For example, when $Au = -\Delta u$, we have

$$\partial_t u(t,x) = \int_0^t \int_{\mathbb{R}^d} k(t-s, x-y) f(s,y)\,\mathrm{d}y\,\mathrm{d}s,$$

where $k(t,x) = \left(\frac{|x|^2}{t^2} - \frac{d}{2t}\right)\frac{e^{-|x|^2/t}}{(2\pi t)^{d/2}}$ is a standard singular kernel of parabolic type:

$$|\partial_t^m \partial_x^\alpha k(t,x)| \le C_{\alpha,m}/(t+|x|^2)^{m+|\alpha|/2}$$

Therefore, by the theory of singular integrals, (1.2) holds for $X = L^p$.

Unfortunately, this argument has not been extended to the settings of a general Banach space X, or more specifically, $X = L^r$ with $r \neq p$. Most existing methods in establishing (1.2) for a general Banach space X rely on the concepts of analytic semigroups [2, 5], operator-valued Fourier multiplier theory and R-boudnedness [9, 10] or H^∞-functional calculus [7].

In this note, we present a simple and fundamental approach to the maximal L^p regularity of parabolic problems, which only uses the concept of singular integrals of Volterra type (which are straightforward modification of the standard singular integrals). Knowledge of analytic semigroups, R-boundedness or H^∞-functional calculus are not required.

2 Singular integrals of Volterra type

Frequently we meet the following type of operators

$$Tf(t) = \int_0^t K(t,s) f(s)\,\mathrm{d}s, \tag{2.3}$$

Notes on maximal L^p regularity

Department of Mathematics, Nanjing University, Nanjing, P.R. China. buyangli@nju.edu.cn

where $f \in L^r(\mathbb{R}_+ \mapsto X)$, $K(t,s)$ is a map from the Banach space X to the Banach space Y for any fixed $0 < s < t < \infty$, and the kernel $K(t,s)$ exhibits singularity such as

$$\|K(t,s)\|_{\mathcal{L}(X,Y)} \le \frac{M}{t-s}, \tag{2.4}$$

$$\|K(t,s) - K(t,s_0)\|_{\mathcal{L}(X,Y)} \le \frac{M|s-s_0|^\sigma}{(t-s_0)^{1+\sigma}} \quad \text{if } t - s_0 \ge 2|s-s_0|, \tag{2.5}$$

$$\|K(t,s) - K(t_0,s)\|_{\mathcal{L}(X,Y)} \le \frac{M|t-t_0|^\sigma}{(t_0-s)^{1+\sigma}} \quad \text{if } t_0 - s \ge 2|t-t_0|, \tag{2.6}$$

for some $\sigma \in (0,1]$, where $\|\cdot\|_{\mathcal{L}(X,Y)}$ denotes the operator norm. A kernel $K(t,s)$ which satisfies the above conditions is called a standard singular kernel of Volterra type. The above properties imply Hörmander's condition: $K(t,s)$ is integrable in any bounded domain $U \times V$ such that $\bar{U} \cap \bar{V} = \emptyset$ and $(t,s) \in U \times V$ implies $t > s > 0$, and

$$\sup_{s,s_0 \in \mathbb{R}_+} \int_{t-s_0 \ge 2|s-s_0|} \|K(t,s) - K(t,s_0)\|_{\mathcal{L}(X,Y)} \, \mathrm{d}t \le M, \tag{2.7}$$

$$\sup_{t,t_0 \in \mathbb{R}_+} \int_{t_0-s \ge 2|t-t_0|} \|K(t,s) - K(t_0,s)\|_{\mathcal{L}(X,Y)} \, \mathrm{d}s \le M. \tag{2.8}$$

A singular integral operator of Volterra type is a bounded linear operator from $L^r(\mathbb{R}_+ \mapsto X)$ to $L^r(\mathbb{R}_+ \mapsto Y)$ for some $r \in (1,\infty]$, given by (2.3) when f has compact support and t does not lie in the support of f, where the kernel $K(t,s)$ satisfies the conditions (2.4)-(2.6). Of course, those conditions also imply Hörmander's conditions (2.7)-(2.8).

Remark 2.1 The above definition mimic the definition of the usual singular integral operators [3, 4, 8]. We should be cautious that a singular kernel of Volterra type does not define the singular integral operator directly. For example, the kernel $K(t,s) = 1/(t-s)$ satisfies (2.4)-(2.6) but the integral operator $Tf(t) = \int_0^t K(t,s)f(s)\,\mathrm{d}s$ is not defined even in the Cauchy principal sense.

The following theorem indicates that singular integral operators of Volterra type preserve the essential properties of standard singular integral operators.

Theorem 2.1 *Let X and Y be reflexive Banach spaces. Then a singular integral operator of Volterra type is of weak-type $(1,1)$ and strong-type (p,p) for $1 < p < \infty$. In particular, if $\|T\|_{L^r(\mathbb{R}_+ \mapsto X) \to L^r(\mathbb{R}_+ \mapsto Y)} = B$, then*

$$\|Tf\|_{L^{1,\infty}(\mathbb{R}_+ \mapsto Y)} \le C(M+B)\|f\|_{L^1(\mathbb{R}_+ \mapsto X)},$$
$$\|Tf\|_{L^p(\mathbb{R}_+ \mapsto Y)} \le C_p(M+B)\|f\|_{L^p(\mathbb{R}_+ \mapsto X)}, \quad 1 < p < \infty.$$

Proof The theorem is proved based on the Calderón-Zygmund decomposition (see Appendix). We only prove the case $1 < r < \infty$, as the case $r = \infty$ can be proved similarly.

Without loss of generality, we can first consider f as a smooth function with compact support and then extend the result to $L^p(\mathbb{R}_+ \mapsto X)$ for $1 \le p < \infty$. Let $f = g + \sum_{Q_j \in \mathcal{Q}} b_j$ be the Calderón–Zygmund decomposition so that both g and b_j are in

$L^1(\mathbb{R}_+ \mapsto X) \cap L^\infty(\mathbb{R}_+ \mapsto X)$ and the sum $\sum_{Q_j \in \mathcal{Q}} b_j$ converges in $L^r(\mathbb{R}_+ \mapsto X)$. Since the operator T is bounded on $L^r(\mathbb{R}_+ \mapsto X)$, it follows that

$$Tf(t) = Tg(t) + \sum_{Q_j \in \mathcal{Q}} Tb_j(t)$$

for almost all $t \in \mathbb{R}_+$. The idea of such decomposition is that, if we let Q_j^* be the unique cube with the same center as Q_j (denoted by s_j), with sides parallel to the sides of Q_j and have side length $l(Q_j^*) = 2l(Q_j)$, then

$$Tb_j(t) = \int_0^t K(t,s)b_j(s)\,\mathrm{d}s = \begin{cases} \int_{Q_j}(K(t,s) - K(t,s_j))b_j(s)\,\mathrm{d}s & \text{for } t \in (Q_j^*)^c \cap \{t > s_j\}, \\ 0 & \text{for } t \in (Q_j^*)^c \cap \{t < s_j\}. \end{cases}$$

Let $b = \sum_{Q_j \in \mathcal{Q}} b_j$ and we note that

$$\begin{aligned} &\big|\{t \in \mathbb{R}_+ : \|Tf(t)\|_Y > 1\}\big| \\ &\le \big|\{t \in \mathbb{R}_+ : \|Tg(t)\|_Y > 1/2\}\big| + \big|\{t \in \mathbb{R}_+ : \|Tb(t)\|_Y > 1/2\}\big| \\ &\le 2^r B^r \|g\|^r_{L^r(\mathbb{R}_+ \mapsto X)} + \big|\{t \notin \cup_j Q_j^* : \|Tb(t)\|_Y > 1/2\}\big| + \big|\cup_j Q_j^*\big| \\ &\le (2^{2r-1}B^r\alpha^{r-1} + C\alpha^{-1})\|f\|_{L^1(\mathbb{R}_+ \mapsto X)} + 2\sum_j \int_{(\cup Q_j^*)^c} \|Tb_j(t)\|_Y\,\mathrm{d}t, \end{aligned}$$

where α is the parameter in the Calderón–Zygmund decomposition. We choose α to satisfy $2^{2r-1}B^r\alpha^{r-1} = 2B$ so that

$$\big|\{t \in \mathbb{R}_+ : \|Tf(t)\|_Y > 1\}\big| \le CB\|f\|_{L^1(\mathbb{R}_+ \mapsto X)} + 2\sum_j \int_{(Q_j^*)^c} \|Tb_j(t)\|_Y\,\mathrm{d}t.$$

For the second term, we have

$$\begin{aligned} &\sum_j \int_{(Q_j^*)^c} \|Tb_j(t)\|_Y\,\mathrm{d}t \\ &= \sum_j \int_{(Q_j^*)^c \cap \{t > s_j\}} \|Tb_j(t)\|_Y\,\mathrm{d}t \\ &\le \sum_j \int_{(Q_j^*)^c \cap \{t > s_j\}} \int_{Q_j} \|K(t,s) - K(t,s_j)\|_{\mathcal{L}(X,Y)} \|b_j(s)\|_X\,\mathrm{d}s\,\mathrm{d}t \\ &\le \sum_j \sup_{s \in Q_j} \int_{t - s_j \ge 2|s - s_j|} \|K(t,s) - K(t,s_j)\|_{\mathcal{L}(X,Y)}\,\mathrm{d}t \int_{Q_j} \|b_j(s)\|_X\,\mathrm{d}s \\ &\le \sum_j M \int_{Q_j} \|b_j(s)\|_X\,\mathrm{d}s \le CM\|f\|_{L^1(\mathbb{R}_+ \mapsto X)}. \end{aligned}$$

Therefore, we have proved the weak-type $(1,1)$ estimate. The strong-type (p,p) estimates for $1 < p < r$ follows from real interpolation.

For $r < p < \infty$, we consider the transpose operator T' defined by $(Tg, f) = (g, T'f)$ for any given $g \in L^r(\mathbb{R}_+ \mapsto X)$ and $f \in L^{r'}(\mathbb{R}_+ \mapsto Y')$. Clearly, the operator T' is

given by

$$T'f(t) = \int_t^\infty K'(t,s)f(s)\,\mathrm{d}s$$

with the kernel $K'(t,s) = K(s,t)' \in \mathcal{L}(Y', X')$. Then T' is bounded from $L^{r'}(\mathbb{R}_+ \mapsto Y')$ to $L^{r'}(\mathbb{R}_+ \mapsto X')$. For $f \in L^\infty(\mathbb{R}_+ \mapsto Y')$ with compact support, we let $f = g + \sum_{Q_j \in \mathcal{Q}} b_j$ be the Calderón–Zygmund decomposition so that both g and b_j are in $L^1(\mathbb{R}_+ \mapsto Y') \cap L^\infty(\mathbb{R}_+ \mapsto Y')$ and the sum $\sum_{Q_j \in \mathcal{Q}} b_j$ converges in $L^{r'}(\mathbb{R}_+ \mapsto Y')$, and

$$T'f(t) = T'g(t) + \sum_{Q_j \in \mathcal{Q}} T'b_j(t)$$

for almost all $t \in \mathbb{R}_+$, and

$$T'b_j(t) = \int_t^\infty K(s,t)'b_j(s)\,\mathrm{d}s = \begin{cases} 0 & \text{for } t \in (Q_j^*)^c \cap \{t > s_j\}, \\ \int_{Q_j} (K(s,t)' - K(s_j,t)')b_j(s)\,\mathrm{d}s & \text{for } t \in (Q_j^*)^c \cap \{t < s_j\}. \end{cases}$$

Then

$$\begin{aligned} &\big|\{t \in \mathbb{R}_+ : \|T'f(t)\|_{X'} > 1\}\big| \\ &\le \big|\{t \in \mathbb{R}_+ : \|T'g(t)\|_{X'} > 1/2\}\big| + \big|\{t \in \mathbb{R}_+ : \|T'b(t)\|_{X'} > 1/2\}\big| \\ &\le 2^r B^r \|g\|^r_{L^r(\mathbb{R}_+ \mapsto Y')} + \big|\{t \notin \cup_j Q_j^* : \|T'b(t)\|_{X'} > 1/2\}\big| + \big|\cup_j Q_j^*\big| \\ &\le (2^{2r-1}B^r\alpha^{r-1} + C\alpha^{-1})\|f\|_{L^1(\mathbb{R}_+ \mapsto Y')} + 2\sum_j \int_{(\cup Q_j^*)^c} \|T'b_j(t)\|_{X'}\,\mathrm{d}t, \end{aligned}$$

where α is the parameter in the Calderón–Zygmund decomposition. We choose α to satisfy $2^{2r-1}B^r\alpha^{r-1} = 2B$ so that

$$\big|\{t \in \mathbb{R}_+ : \|T'f(t)\|_{X'} > 1\}\big| \le CB\|f\|_{L^1(\mathbb{R}_+ \mapsto Y')} + 2\sum_j \int_{(Q_j^*)^c} \|T'b_j(t)\|_{X'}\,\mathrm{d}t.$$

For the second term, we have

$$\begin{aligned} &\sum_j \int_{(Q_j^*)^c} \|T'b_j(t)\|_{X'}\,\mathrm{d}t \\ &= \sum_j \int_{(Q_j^*)^c \cap \{t < s_j\}} \|T'b_j(t)\|_{X'}\,\mathrm{d}t \\ &\le \sum_j \int_{(Q_j^*)^c \cap \{t < s_j\}} \int_{Q_j} \|K(s,t)' - K(s_j,t)'\|_{\mathcal{L}(Y',X')} \|b_j(s)\|_{Y'}\,\mathrm{d}s\,\mathrm{d}t \\ &\le \sum_j \sup_{s \in Q_j} \int_{s_j - t \ge 2|s - s_j|} \|K(s,t) - K(s_j,t)\|_{\mathcal{L}(X,Y)}\,\mathrm{d}t \int_{Q_j} \|b_j(s)\|_{Y'}\,\mathrm{d}s \\ &\le \sum_j M \int_{Q_j} \|b_j(s)\|_{Y'}\,\mathrm{d}s \le CM\|f\|_{L^1(\mathbb{R}_+ \mapsto Y')}. \end{aligned}$$

Therefore, we have proved the weak-type $(1,1)$ estimate. The strong-type (p', p') estimates for $1 < p' < r'$ follows from real interpolation. In other words, T' is bounded

from $L^{p'}(\mathbb{R}_+ \mapsto Y')$ to $L^{p'}(\mathbb{R}_+ \mapsto X')$ for $1 < p' < r'$. By a simple duality argument, this implies that T is bounded from $L^p(\mathbb{R}_+ \mapsto X)$ to $L^p(\mathbb{R}_+ \mapsto Y)$ for $r < p < \infty$.

Overall, T is bounded from $L^p(\mathbb{R}_+ \mapsto X)$ to $L^p(\mathbb{R}_+ \mapsto Y)$ for $1 < p < \infty$. □

3 Maximal L^p regularity of parabolic equations

Consider the parabolic problem

$$\partial_t u - \sum_{i,j=1}^{d} \partial_j(a_{ij}(x)\partial_i u) + \sum_{j=1}^{d} b_j(x)\partial_j u + c(x)u = f \tag{3.1}$$

with the initial condition $u(0,x) \equiv 0$, where the coefficients a_{ij}, b_j and c are bounded, measurable and satisfying the strongly ellipticity condition:

$$\Lambda^{-1}|\xi|^2 \le \sum_{i,j=1}^{d} a_{ij}(x)\xi_i\xi_j \le \Lambda|\xi|^2, \quad \forall\, x, \xi \in \mathbb{R}^d.$$

This corresponds to (3.1) with $Au = -\sum_{i,j=1}^{d} \partial_j(a_{ij}(x)\partial_i u) + \sum_{j=1}^{d} b_j(x)\partial_j u + c(x)u$.

Let $G(t,x,y)$ denote the Green function of the parabolic equation so that the solution of (3.1) is given by

$$u(t,x) = \int_0^t \int_{\mathbb{R}^d} G(t-s,x,y) f(s,y)\,\mathrm{d}y\,\mathrm{d}s.$$

Thus

$$\partial_t u(t,x) = f(t,x) + \int_0^t \int_{\mathbb{R}^d} \partial_t G(t-s,x,y) f(s,y)\,\mathrm{d}y\,\mathrm{d}s.$$

Let $X = L^r$ and let the mapping from $f \in L^r(\mathbb{R}_+; X)$ to $\partial_t u - f \in L^r(\mathbb{R}_+; X)$ be denoted by T. Then $Tf(t) = \int_0^t K(t-s)f(s)\,\mathrm{d}s$ when f has compact support and t is not in the support of f, where the operator-valued kernel $K(t) : L^r \to L^r$ can be expressed as

$$[K(t)g](x) = \int_{\mathbb{R}^d} \partial_t G(t,x,y) g(y)\,\mathrm{d}y.$$

The kernel $K(t)$ obeys the standard estimates:

$$\|K(t-s)\|_{\mathcal{L}(L^r,L^r)} \le \frac{C}{t-s}, \qquad \text{for } 0 < s < t < \infty, \tag{3.2}$$

$$\|\partial_t K(t-s)\|_{\mathcal{L}(L^r,L^r)} \le \frac{C}{(t-s)^2}, \quad \text{for } 0 < s < t < \infty, \tag{3.3}$$

which indicate that $T : L^r(\mathbb{R}_+; X) \to L^r(\mathbb{R}_+; X)$ is a singular integral operator of Volterra type. By Theorem 2.1, this operator must be bounded on $L^p(\mathbb{R}_+; X)$ for all $1 < p < \infty$.

Remark 3.1 By this approach, the maximal regularity

$$\|\partial_t u\|_{L^p(\mathbb{R}_+;L^r)} + \|Au\|_{L^p(\mathbb{R}_+;L^r)} \le C\|f\|_{L^p(\mathbb{R}_+;L^r)}, \quad \text{for } 1 < p, r < \infty$$

reduces to the homogeneous estimate

$$\|\partial_t u\|_{L^r(\mathbb{R}_+;L^r)} + \|Au\|_{L^r(\mathbb{R}_+;L^r)} \le C\|f\|_{L^r(\mathbb{R}_+;L^r)}, \quad \text{for } 1 < r < \infty.$$

This approach can also be applied to problems defined on a finite time interval as well as problems defined on the bounded domain.

Appendix: Calderón–Zygmund decomposition on $\mathbb{R}_+$

Let $\mathbb{Z} = \{0, \pm 1, \pm 2 \cdots\}$ denote the set of all integers and let $\mathbb{N} = \{0, 1, 2, \cdots\}$ denote the set of all natural numbers. For any integers $n \in \mathbb{Z}$ and $k \in \mathbb{N}$ we define $Q_{n,k} = (2^n k, 2^n(k+1)]$, called a dyadic cube. Then $\mathscr{Q}_+ := \{Q_{n,k} : n \in \mathbb{Z} \text{ and } k \in \mathbb{N}\}$ is called the set of all dyadic cubes on the half-line $\mathbb{R}_+$. For any two dyadic cubes $Q_1, Q_2 \in \mathscr{Q}_+$, either $Q_1 \subset Q_2$ or $Q_2 \subset Q_1$ or $Q_1 \cap Q_2 = \emptyset$. This is often referred to as the nesting property of dyadic cubes.

Proposition *Let X be a Banach space, $f \in L^1(\mathbb{R}_+ \mapsto X)$ and $\alpha > 0$. Then there is a decomposition*

$$f = g + \sum_{j=1}^{\infty} b_j$$

which satisfies that

(1) $\|g\|_{L^1(\mathbb{R}_+\mapsto X)} \le \|f\|_{L^1(\mathbb{R}_+\mapsto X)}$ *and* $\|g\|_{L^\infty(\mathbb{R}_+\mapsto X)} \le 2\alpha$,

(2) *the functions b_j are supported in disjoint dyadic cubes $Q_j \in \mathcal{Q}_+$, respectively,*

(3) $\int_{Q_j} b_j(t)\,\mathrm{d}t = 0$, $\int_{Q_j} \|b_j\|_X\,\mathrm{d}t \le 4\alpha|Q_j|$,

(4) $\sum_j |Q_j| \le \|f\|_{L^1(\mathbb{R}_+\mapsto X)}/\alpha$,

(5) *if $f \in L^\infty(\mathbb{R}_+ \mapsto X)$ with compact support in $\mathbb{R}_+$, then $g \in L^\infty(\mathbb{R}_+ \mapsto X)$ with compact support in $\mathbb{R}_+$, $\int_{\mathbb{R}_+} \|\sum_{j=k}^{l} b_j\|_X^r\,\mathrm{d}t \le 2^r \int_{\cup_{j=k}^{l} Q_j} \|f\|_X^r\,\mathrm{d}t$ and so the series $\sum_{j=1}^{\infty} b_j$ converges in $L^r(\mathbb{R}_+ \mapsto X)$ for any $1 \le r < \infty$.*

Proof Let us say that a dyadic cube Q is bad if $\frac{1}{|Q|}\int_Q \|f(t)\|_X\,\mathrm{d}t > \alpha$, and good otherwise. A maximal bad dyadic cube is a bad dyadic cube such that any dyadic cube strictly containing it is good. Since $f \in L^1(\mathbb{R}_+ \mapsto X)$, any bad dyadic cube is contained in a maximal bad dyadic cube. Let $\mathcal{Q}$ be the collection of all maximal bad dyadic cubes. By the nesting property of dyadic cubes, cubes in $\mathcal{Q}$ are disjoint. For any $Q \in \mathcal{Q}$, $\frac{1}{|Q|}\int_Q \|f(t)\|_X\,\mathrm{d}t > \alpha$ and $\frac{1}{|Q'|}\int_{Q'} \|f(t)\|_X\,\mathrm{d}t \le \alpha$ for any dyadic cube Q' strictly containing Q. Therefore,

$$\alpha < \frac{1}{|Q|}\int_Q \|f(t)\|_X\,\mathrm{d}t \le 2\alpha.$$

For any dyadic cube Q outside $\cup\mathcal{Q}$, $\frac{1}{|Q|}\int_Q \|f(t)\|_X\,\mathrm{d}t \le \alpha$. By the Lebesgue differentiation theorem,

$$\|f(t)\|_X = \lim_{\mathrm{diam}(Q)\to 0} \frac{1}{|Q|}\int_Q \|f(s)\|_X\,\mathrm{d}s \le 2\alpha$$

for almost all t outside $\cup\mathcal{Q}$, where Q extends over all sequence of dyadic cubes disjoint from $\cup\mathcal{Q}$ and containing t.

Let

$$b_j(t) = \left(f(t) - \frac{1}{|Q_j|} \int_{Q_j} f(s)\,\mathrm{d}s \right) 1_{Q_j}(t), \qquad g(t) = \frac{1}{|Q_j|} \int_{Q_j} f(s)\,\mathrm{d}s$$

for $t \in Q_j \subset \mathcal{Q}$. Let $g = f$ outside $\cup\mathcal{Q}$. Then $f = g + \sum_{Q_j \in \mathcal{Q}} b_j$ satisfies the requirements. □

References

[1] Calderón, *Singular integrals*, Colloquium Lectures given in Aug 31–Sep 3, 1965 at the Seventieth Summer Meeting of the American Mathematical Society held in Ithaca, New York.

[2] P. Cannarsa and V. Vespri, *On maximal L^p regularity for the abstract Cauchy problem*, Boll. Un. Mat. Ital. B, 5 (1986), pp. 165-175.

[3] L. Grafakos, *Classical Fourier analysis*, Springer Science+Business Media, LLC, 2008.

[4] L. Grafakos, *Modern Fourier analysis*, Springer Science+Business Media, LLC, 2009.

[5] M. Hieber and J. Prüss, *Heat kernels and maximal L^p-L^q estimates for parabolic evolution equations*, Comm. Partial Differential Equations, 22 (1997), pp. 1647-1669.

[6] B.F. Jones, *A Class of Singular Integrals*, American J. Math., 86 (1964), pp. 441-462.

[7] C.L. Merdy, *H^∞-functional calculus and applications to maximal regularity*, Publ. Math. UFR Sci. Tech. Besancon. 16 (1998), pp. 41-77.

[8] E.M. Stein, *Singular integrals and differentiability properties of functions*, Princeton University Press, 1970.

[9] L. Weis, *A new approach to maximal L^p-regularity*, Evolution equations and their applications in physical and life sciences (Bad Herrenalb, 1998), Dekker, New York, 2001, pp. 195-214.

[10] L. Weis, *Operator-valued Fourier multiplier theorems and maximal L^p-regularity*, Math.Ann., 319 (2001), pp.735-758.